

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/348180996>

BIOMARKERS OF MYOCARDIAL FIBROSIS AND RENAL DYSFUNCTION IN HYPERTENSIVE PATIENTS WITH TYPE 2 DIABETES MELLITUS

Article in *Pharmacophore* · December 2020

CITATIONS

0

READS

42

7 authors, including:



Alexander E Berezin

Zaporozhye State Medical University

697 PUBLICATIONS 2,206 CITATIONS

SEE PROFILE


Some of the authors of this publication are also working on these related projects:



Impaired immune phenotype of circulating endothelial cell-derived microparticles as novel biomarker for diabetes mellitus [View project](#)



Brazilian Network of Research in Meta-analysis - BRAMETIS [View project](#)



BIOMARKERS OF MYOCARDIAL FIBROSIS AND RENAL DYSFUNCTION IN HYPERTENSIVE PATIENTS WITH TYPE 2 DIABETES MELLITUS

Oleksandr M Bilovol¹, Iryna I. Kniazkova², Oleksandr M. Kirienko³, Maryna V Bogun⁴, Denis M. Kirienko⁵, Lilya P. Abramova⁶, Alexander E Berezin^{7*}

1. *Member of the Presidium of the Academy of Medical Sciences of Ukraine, MD, PhD, Professor of Clinical Pharmacology and Internal Medicine Department, Kharkiv National Medical University, Kharkiv, Ukraine.*
2. *Professor, MD, PhD, Head of Clinical Pharmacology and Internal Medicine department, Kharkiv National Medical University, Kharkiv, Ukraine.*
3. *MD, assistant professor of Clinical Pharmacology and Internal Medicine department, Kharkiv National Medical University, Kharkiv, Ukraine.*
4. *MD, PhD, associate professor, statistic specialist, Kharkiv National Medical University, Kharkiv, Ukraine.*
5. *MD, PhD, associate professor of Clinical Pharmacology and Internal Medicine department, Kharkiv National Medical University, Kharkiv, Ukraine.*
6. *Senior Researcher, Kharkiv National Medical University, Kharkiv, Ukraine.*
7. *Professor, MD, PhD, Senior Consultant of Therapeutic Unit, Internal Medicine Department, State Medical University of Zaporozhye, 26, Mayakovsky av., Zaporozhye, 69035, Ukraine.*

ARTICLE INFO

Received:

07 Oct 2020

Received in revised form:

14 Dec 2020

Accepted:

18 Dec 2020

Available online:

28 Dec 2020

Keywords: hypertension, type 2 diabetes mellitus, soluble suppressor tumorigenicity-2, neutrophilic gelatinase-associated lipocalin, diastolic function, diastolic function.

ABSTRACT

The aim of the study was to investigate the interaction of circulating levels of sST2, neutrophilic gelatinase-associated lipocalin, and cystatin C with cardiac remodeling in hypertensive patients depending on the presence of T2DM. **Methods:** The study included 76 mild-to-moderate hypertensive patients, of whom 48 patients were with stage II grade 2 of hypertension and T2DM, and 28 patients with stage II of hypertension without T2DM. The control group consisted of 20 healthy volunteers (mean age of 54.5 ± 3.1 years). All examined persons underwent general clinical examination, determination of indicators of carbohydrate metabolism and lipid profile; insulin resistance was assessed using the HOMA-IR index. Serum levels of soluble suppressor tumorigenicity-2 (sST2), cystatin C, and NGAL were measured using kits for enzyme-linked immunosorbent assay. The structural and functional parameters of the heart and the diastolic function of the left ventricle were analyzed using echocardiography with Doppler sonography. **Results:** It was found that concentric and eccentric LV hypertrophy, and diastolic dysfunction, occurred more often in hypertensive patients with T2DM than in the hypertensive patients without T2DM. In T2DM patients eccentric and concentric LV remodeling was associated with the levels of sST2 ($r = 0.53$; $P = 0.001$ and $r = 0.55$; $P = 0.001$, respectively), cystatin C ($r = 0.40$; $P = 0.001$ and $r = 0.44$; $P = 0.001$, respectively) and NGAL ($r = 0.38$; $P = 0.001$ and $r = 0.44$; $P = 0.001$, respectively). Backward stepwise regression analyses showed that the most influential determinants of depending variable (LV remodeling) in T2DM patients were sST2 (B-coefficient = 1.9, SD = 0.03; $p = 0.001$) and cystatin C (B-coefficient = 0.92, SD = 0.03; $p = 0.022$), whereas NGAL exhibited weak discriminative ability (B-coefficient = 0.21, SD = 0.01; $p = 0.16$). **Conclusion:** Cardiac remodeling in hypertensive patients having T2DM was significantly associated with higher levels of sST2 and cystatin C. sST2 had the best potency to predict LV remodeling when compared with NGAL and other biomarkers.

Copyright © 2013 - All Rights Reserved – Pharmacophore

To Cite This Article: Oleksandr M Bilovol, Iryna I. Kniazkova, Oleksandr M. Kirienko, Maryna V Bogun, Denis M. Kirienko, Lilya P. Abramova, (2020), "Biomarkers of myocardial fibrosis and renal dysfunction in hypertensive patients with type 2 diabetes mellitus", *Pharmacophore*, 11(6), 132-140.

Corresponding Author: Alexander E. Berezin; Professor, MD, PhD, Senior Consultant of Therapeutic Unit, Internal Medicine Department, State Medical University of Zaporozhye, 26, Mayakovsky av., Zaporozhye, 69035, Ukraine. Email: aeberezin@gmail.com

Introduction

Arterial hypertension (AH) is a disease that occurs when there is a change in cardiac output, regional vascular resistance, or both resulting in a constant increase in blood pressure of the arteries of individuals [1]. Diabetes mellitus is a common endocrine disorder caused by the reduction of insulin production through the pancreas, which results in an increase in glucose concentrations in the blood [2]. One of the important medical and social problems of our time is the burden of comorbidity including a combination of arterial hypertension and type 2 diabetes mellitus (T2DM), which causes an increase in the risk of premature mortality [3]. Hemodynamic disorders together with metabolic disorders in patients with AH and T2DM lead to damage in target organs and early development of cardiovascular (CV) complications with both macro-angiopathy and micro-angiopathy [4]. It is known that unfavorable changes in the heart and kidneys, in turn, significantly worsen the prognosis of patients with AH and T2DM increasing the risk of CV complications [5]. Therefore, the study of target organ damage in patients having both AH and type 2 DM is important for improvement in the prognosis and point-of-care strategy of the patients [6].

Adverse cardiac remodeling in hypertension is caused by a complex interaction between genetic, neurohormonal, and inflammatory changes affecting cardiac myocytes and extracellular matrix [7-9]. In this regard, the increasing clinical significance is assigned to the biomarkers of myocardial stress, dysfunction, and cellular damage [10]. Among the biomarkers of myocardial fibrosis, soluble suppressor of tumorigenicity-2 (sST2) is actively studied and exhibits conflicting results in the prediction of poor prognosis in patients having T2DM [11]. However, there is no strong agreement on the prognostic value of modern indicators of renal damage and cystatin C dysfunction, and neutrophilic gelatinase-associated lipocalin (NGAL) [12, 13]. The study of the relationship of these biomarkers with clinical, instrumental, and other biochemical parameters in hypertension in combination with T2DM seems to be a promising direction in the management strategy of this cohort of patients.

The aim of the study was to investigate the interaction of circulating levels of sST2, neutrophilic gelatinase-associated lipocalin, and cystatin C with cardiac remodeling in hypertensive patients depending on the presence of T2DM.

Methods

The single-center cross-sectional study included 76 hypertensive patients, 48 patients (29 males and 19 females, mean age of 54.3 ± 3.2 years) had T2DM, and 28 patients (18 males and 10 females, mean age of 54.8 ± 3.5 years) did not have T2DM. All patients were examined in accordance with the recommendations of the European Society of Hypertension and the European Society of Cardiology (ESH/ESC, 2018) [14, 15] and the American Diabetes Association (ADA, 2017) [16].

The exclusion criteria were the following: stage III, grade 3 of hypertension, history of exertional angina, history of myocardial infarction and transient ischemic attack/stroke, III/IV NYHA functional class of chronic heart failure, valvular heart disease, connective tissue diseases, malignancy, thyroid dysfunction, type 1 DM, patients undergoing insulin therapy; reduced glomerular filtration rate < 60 mL/min; and left ventricular ejection fraction $< 50\%$.

All patients underwent antihypertensive therapy (a combination of generally accepted drug classes: ACE inhibitors, calcium channel antagonists, beta-blocker [carvedilol] in optimal therapeutic doses). All patients received rosuvastatin at a dose of 20 mg and acetylsalicylic acid at a daily dose of 75 mg. A combination of metformin and gliclazide was prescribed to improve hyperglycemia. The control group consisted of 20 healthy individuals (mean age of 54.5 ± 3.1 years).

Ethical declaration

Informed consent was obtained from all patients for participation in the study.

General anthropometric, clinical, and physical examinations

Conventional anthropometric measurements (height, weight, waist circumference, hip-to-waist ratio, body mass index [BMI]) were performed. Clinical history was recorded and specified. All patients underwent general clinical and physical examination, office blood pressure (BP) and heart rate (HR) measure, as well as ambulatory BP monitoring at baseline.

Echocardiography

Echocardiography was carried out on the State Institution "National Institute of Therapy named after L. T. Malaya of NAMS of Ukraine". We used ultrasound machine "GE Medical Systems" (Germany) by a phase sensor with a modulated frequency of 2.25-3 MHz in M- and B-modes in accordance with the recommendations of the American Society of Echocardiography (2015) [17]. LV myocardial mass (LVMM) was calculated and the LVMM index (LVMMI) was determined as the ratio of LVMM to body surface area. The types of LV remodeling (eccentric LVH, concentric LVH, concentric remodeling) were determined in accordance with the ASE guidelines (2015) [17].

The analysis of LV diastolic function was carried out during the registration of the transmitral diastolic flow in the pulse-wave Doppler mode [18]. The maximum rate of late filling of the LV in tissue mode (A), the maximum rate of late filling of the LV in tissue mode (a), the maximum rate of the early atrial filling (E), the maximum rate of early filling of the LV in tissue mode (e), the ratio of the rates of early and late fillings (E/A), the ratio of the maximum rates of early and late LV fillings in the

tissue mode (e/a), the time of deceleration of the rate of early diastolic flow (DT), isovolumic relaxation time (IVRT) were also determined.

Glomerular filtration rate measurement

Calculation of glomerular filtration rate (GFR) was calculated by CKD-EPI formula [19]

Insulin resistance determination

Insulin resistance was calculated as Homeostatic Assessment Model of Insulin Resistance (HOMA-IR) using the appropriate equation [20]:

$$\text{HOMA-IR} = \text{fasting insulin (mU/L)} \times \text{fasting glucose (mmol/L)} / 22.5$$

Biomarkers measurement

Blood samples were drawn in the morning following overnight fasting (at 7-8 a.m.) into barcoded silicone test tubes. Then samples were centrifuged upon permanent cooling at 6,000 rpm for 3 minutes and then plasma was collected to be immediately refrigerated. Each aliquot was stored at -70°C.

Fasting blood serum glucose, glycosylated hemoglobin (HbA1c) levels in whole blood, insulin, and lipid profile parameters were determined using commercial kits manufactured by DRG (USA) using Labline-90 analyzer (Austria)

The concentration of sST2 ("Critical Diagnostics", USA), cystatin C (Bio Vendor, Czech Republic), and NGAL ("Hycult Biotech", Netherlands) were measured in serum with commercial kits for enzyme-linked immunosorbent assay.

Statistics

Computer processing of the study results was made using the Statistica 8.0 software package (StatSoft Inc, USA). The mean (M), variance, standard deviation, median (m), significance, and significance level (p) were calculated. For assessment of the relationship between the indicators, the method of correlation analysis was used with the calculation of Pearson's correlation coefficients (with a normal distribution) and Spearman's correlation coefficients (with a distribution that differs from normal). Backward stepwise multiple regression analysis was performed to establish the most significant determinants of LV remodeling entering the following variables: age, sex, waist circumference, WHR, BMI, CV risk factors, HOMA-IR, and biomarkers (sST2, NGAL, cystatin C). Differences were considered significant at the level of statistical significance $p < 0.05$.

Results

Basic demographic, clinical, and anthropometric characteristics of the patient study population are reported in Table 1. There were no significant differences between both populations of hypertensive patients with and without T2DM.

Table 1: Basic demographic, clinical and anthropometric characteristics of patient study population

Variables	Healthy volunteers (n=20)	Entire patient cohort (n=76)	Hypertensive patients (n=76)		P- value
			T2DM patients (n=48)	Non-T2DM patients (n=28)	
Age, year	54.5 (50-57)	54.5 (50-58)	54.3 (51-57)	54.8 (50-58)	NS
Male, n (%)	12 (60)	47 (61.8)	29 (60.4)	18 (64.3)	NS
Dyslipidemia, n (%)	-	73 (96.1)#	47 (97.9)	26 (96.4)	NS
Smoking, n (%)	9 (45)	35 (46.1)	22 (45.8)	13 (46.4)	NS
Abdominal obesity, n (%)	-	30 (39.5)#	19 (39.6)	11 (39.3)	NS
Microalbuminuria, n (%)	-	22 (28.9)#	14 (29.2)	8 (28.6)	NS
LV hypertrophy, n (%)	-	55 (72.4)#	40 (83.3)	15 (53.6)	0.001
BMI, kg/m ²	22.13 ± 0.35	25.26 ± 2.47#	25.63 ± 2.45	24.91 ± 2.56	NS
Waist circumference, sm	78.9±2.73	83.0±3.61	84.1±3.67	82.3±3.58	NS
WHR, units	0.78±0.02	0.85±0.04	0.86±0.05	0.84±0.04	NS

Notes: data are given as mean ± SD and median, # - significant difference between healthy volunteers and entire patient' cohort (P1-2<0.05).

Abbreviations: WHR, Waist-to-hip ratio; BMI, body mass index; NS, not significant; SD, standard deviation.

Analysis of echocardiographic data showed that the most pronounced disorders of the structural and functional state of the myocardium were found in hypertensive patients with T2DM. LV hypertrophy was detected in 40 (83.3% of patients with T2DM and in 15 (53.56%) patients without T2DM (Table 2). Moreover, in patients with T2DM, unfavorable types of myocardial remodeling - concentric and eccentric LV hypertrophy (in 64.58% and 18.75%, respectively) were revealed, whereas in patients without type 2 DM, concentric remodeling and concentric LV hypertrophy prevailed (in 17.85% and 35.71%, respectively).

Table 2: Hemodynamics and biomarkers in patient study population

Variables	Healthy volunteers (n=20)	Entire patient cohort (n=76)	Hypertensive patients (n=76)		P- value
			T2DM patients (n=48)	Non-T2DM patients (n=28)	
Office SBP, mm Hg	124 ± 4	135 ± 5	134 ± 5	136 ± 6	NS
Office DBP, mm Hg	76 ± 5	79 ± 5	80 ± 4	78 ± 5	NS
LVEDV, mL	93 ± 2	113 ± 3#	117 ± 4	108 ± 3	NS
LVESV, mL	30 ± 2	42 ± 2#	43 ± 3	39 ± 3	0.05
LVEF, %	64 ± 3	62 ± 1#	62 ± 3	62 ± 3	NS
LVMMI, g/m ²	82.3 ± 0.07	133.45 ± 1.93#	141.35 ± 2.01	123.56 ± 1.83	0.05
E/e'	5.31 ± 0.08	6.01 ± 0.12#	6.30 ± 0.15	5.73 ± 0.13	0.05
GFR, mL/min/1.73 m ²	102 ± 7.63	93 ± 5.63	87 ± 5.78	95 ± 5.59	NS
HOMA-IR	1.66 ± 0.58	5.31 ± 2.87#	5.63 ± 3.48	4.49 ± 2.51	NS
Fasting glucose, mmol/L	4.27 ± 1.35	5.73 ± 1.54#	6.14 ± 1.76	5.43 ± 1.32	NS
HbA1c, %	4.38 ± 1.51	6.17 ± 1.37#	6.59 ± 1.59	5.58 ± 1.39	NS
TC, mmol/L	5.2 ± 1.13	6.23 ± 1.13#	6.35 ± 1.17	5.91 ± 1.15	0.05
HDL-C, mmol/L	1.3 ± 0.12	1.00 ± 0.12#	0.93 ± 0.15	1.08 ± 0.09	NS
LDL-C, mmol/L	3.05 ± 0.05	4.31 ± 0.07#	5.03 ± 0.06	3.95 ± 0.08	0.045
TG, mmol/L	1.7 ± 0.05	2.23 ± 0.04#	2.28 ± 0.09	2.05 ± 0.08	NS
sST2, ng/mL	19.1 (16.19-26.12)	31.29 (24.32 - 42.35)#	35.29 (31.33 - 42.35)	27.15 (24.32 - 33.18)	0.001
NGAL, ng/mL	9.73 (6.83 - 14.12)	15.19 (12.0 - 20.19) #	17.12 (13.15 - 20.19)	14.26 (12.0 - 17.25)	0.058
Cystatin C, ng/mL	82.0 (68.79 - 108.12)	193.0 (101.40-302.10)#	269.0 (179.01- 328.13)	106.0 (98.93 - 270.12)	0.001

Notes: data are given as mean ± SD and median, # - significant difference between healthy volunteers and entire patient' cohort (P1-2<0.05).

Abbreviations: SBP – systolic blood pressure; DBP – systolic blood pressure; HbA1c, glycosylated hemoglobin; GFR, glomerular filtration rate, TC, total cholesterol; LDL, low-density lipoproteins; HDL, high-density lipoproteins; TG, triglycerides.

Hypertensive patients with T2DM had the highest LV myocardial mass index (LVMI), end-diastolic and end-systolic LV dimensions (EDD and ESD, respectively), end-diastolic and end-systolic LV volumes (EDV and ESV, respectively) in comparison with the group of patients with AH. The mean values of LVMM index in patients having T2DM were significantly higher than in patients without T2DM and the control group (p <0.05). The differences in all presented echocardiographic parameters were statistically significant between hypertensive patients with and without T2DM.

LV diastolic filling abnormalities were also found in hypertensive patients with T2DM and without it. However, both early (E) and late (A) diastolic filling velocities were more significantly reduced in T2DM patients than non-T2DM individuals and healthy volunteers. However, 16.7% of patients with T2DM and 35.7% of patients without T2DM did not have diastolic filling abnormalities.

In 52 (68.4%) patients who took part in the study, a slight decrease of renal function was diagnosed (GFR = 60-89 ml/min/1.73 m²). Normal renal function (GFR ≥90 ml/min/1.73 m²) was determined in 24 (31.6%) patients. Pairwise comparison of the groups revealed a statistically significant predominance of the reduced renal function in the group of patients with T2DM in comparison to those without it (p <0.001). The lowest estimated GFR was found in patients of group 1 in comparison with the patients of group 2 (p <0.01).

A comparative analysis of the levels of sST2, NGAL, and cystatin C in patient study populations is reported in Figure 1. In fact, the levels of sST2, NGAL, and cystatin C were significantly higher in T2DM patients when compared with non-T2DM individuals and healthy volunteers.




Figure 1A




Figure 1B




Figure 1C

Figure 1: Circulating levels of biomarkers in patients with and without T2DM in comparison with healthy volunteers.
Figure 1A: sST2; Figure 1B: Cystatin C; Figure 1C: NGAL

Abbreviations: NS, not significant. The lines indicate comparisons between variables.

The concentrations of the biomarkers (sST2, NGAL, and cystatin C) in patients with T2DM did not differ significantly according to gender, overweight, or obesity (for all $p=0.46$).

Correlations between HOMA-IR, anthropometric parameters, age, LV hypertrophy, diastolic filling abnormality, and biomarkers

In entire T2DM population, HOMA-IR was significantly and positively correlated with serum TG ($r=0.48$; $p=0.002$), WHR ($r=0.50$; $p=0.001$), waist circumference ($r=0.48$; $p=0.001$), LH hypertrophy ($r=0.46$; $p=0.001$), sST2 ($r=0.44$; $p=0.042$), BMI ($r=0.42$; $p=0.002$), age ($r=0.44$; $p=0.012$), diastolic filling abnormality ($r=0.42$; $p=0.026$), microalbuminuria ($r=0.32$; $P=0.002$), and smoking ($r=0.24$; $P=0.001$), while inverse correlation of HOMA-IR with HDL-C ($r=-0.40$; $p=0.02$) was observed.

Serum levels of NGAL and cystatin C were positively correlated with microalbuminuria ($r=0.52$; $P=0.001$ and $r=0.54$; $P=0.002$, respectively), WHR ($r=0.44$; $p=0.001$ and $r=0.43$; $P=0.043$, respectively), waist circumference ($r=0.42$; $p=0.022$ and $r=0.44$; $P=0.022$, respectively), LH hypertrophy ($r=0.42$; $p=0.026$ and $r=0.42$; $P=0.016$, respectively).

Correlations between eccentric and concentric forms of LV hypertrophy and biomarkers' levels

We found positive correlation between eccentric and concentric LV remodeling and levels of sST2 ($r=0.50$; $P=0.042$ and $r=0.54$; $P=0.001$, respectively), WHR ($r=0.42$; $P=0.022$ and $r=0.44$; $P=0.001$, respectively). The levels of cystatin C ($r=0.28$; $P=0.042$) and NGAL ($r=0.26$; $P=0.022$) exhibited weak association with concentric LV remodeling in entire hypertensive patient population and did not correlated with eccentric LV remodeling.

Correlations between eccentric and concentric forms of LV hypertrophy and biomarkers' levels in T2DM subpopulation

In T2DM patients eccentric and concentric LV remodeling was associated with the levels of sST2 ($r=0.53$; $P=0.001$ and $r=0.55$; $P=0.001$, respectively), cystatin C ($r=0.40$; $P=0.001$ and $r=0.44$; $P=0.001$, respectively) and NGAL ($r=0.38$; $P=0.001$ and $r=0.44$; $P=0.001$, respectively).

Backward Wald binary logistic regression analysis on factors associated with LV remodeling in T2DM patients

Backward stepwise regression analyses showed that the most influential determinants of depending variable (LV remodeling) in T2DM patients were sST2 (B-coefficient = 1.9, SD = 0.03; $p=0.001$) and cystatin C (B-coefficient = 0.92, SD = 0.03; $p=0.022$), whereas NGAL exhibited weak discriminative ability (B-coefficient = 0.21, SD = 0.01; $p=0.16$). Other variables, such as WHR, BMI, a number of CV risk factors, HOMA-IR, had insufficient predictive potency in both unadjusted and adjusted for BMI regression models.

Discussion

The results of our study showed that the levels of sST2 and NGAL were significantly higher in patients with T2DM with mild renal dysfunction, whereas, there were no significant differences in the levels of cystatin C in both subpopulations of the patients. Nevertheless, eccentric and concentric LV remodeling was closely associated with the levels of sST2, cystatin C, and NGAL in patients having T2DM only.

We suggest that these biomarkers can be promising predictors for LV remodeling in hypertensive patients having T2DM and that sST2 is a potential target for therapy among these individuals. Recently other investigators gave their comments regarding use in routine clinical practice biomarkers of fibrosis (sST2) and kidney function (cystatin C and NGAL) to predict the natural evolution of both cardiac dysfunction and kidney dysfunction [21-24]. In this context, sST2 appeared to be better than others because it independently predicted CV events and diseases from early stages of chronic kidney disease to the late stage of one [25, 26]. However, there are conflicting results reflecting serious difficulties to extrapolate the predictive ability of these biomarkers to T2DM patients at different stages of chronic kidney disease [24]. Such biomarkers as NGAL, cystatin C, and KIM-1 were found to be powerful predictors for declining kidney function renal events and CV complications in patients with moderate and severe chronic kidney disease [27-30]. We firstly obtained data that clarify the role of sST2 in relation to LV remodeling among T2DM patients not having heart failure and declining GFR. Perhaps, sST2 is desirably considered as an early biomarker of adverse cardiac remodeling in the diabetic population. Therefore, normal and near-normal GFR with microalbuminuria and elevation of circulating levels of NGAL reflecting early stage of diabetes mellitus-induced nephropathy do not have pervasive predictive potency for adverse cardiac remodeling in hypertensive patients with T2DM [31, 32].

Recent clinical studies and prospective ASIAN-HF (Asian Sudden Cardiac Death in Heart Failure) Registry have revealed that in both heart failure (HF) phenotypes (HF with reduced [HFrEF] and preserved [HFpEF] ejection fraction) T2DM (versus non-T2DM) was associated with smaller indexed LV diastolic volumes and higher mitral E/e' ratio that indicated on diastolic filling abnormality [33-35]. There was a predominance of eccentric hypertrophy in HFrEF and concentric hypertrophy in HFpEF in patients having T2DM. In this context, biomarkers of adverse cardiac remodeling, fibrosis, and biomechanical stress, such as sST2, appear to be attractive for further evaluation. The different ability of sST2 and cystatin C to be associated with

LV remodeling and diastolic filling abnormality in T2DM and non-T2DM hypertensive patients, which was established in our study, requires performing new investigations in large clinical studies.

We suggest that metabolic alterations of intermediate substrate metabolism and oxidative stress rather than direct tubular kidney damage per se account for maladaptive cardiac remodeling and diastolic filling abnormality. Despite worsening kidney function can negatively impact cardiac structure and function through increased pressure overload and induce LV hypertrophy and consequently the development of HF, extracellular accumulation of collagen remains the primary mechanism for the occurrence of diastolic dysfunction in hypertensive patients with T2DM [36]. This sST2 is a mostly preferable biomarker that is suggesting higher susceptibility to the prediction of deleterious effects of chronic pressure overload due to hypertension and chronic kidney disease, among this T2DM population.

Study limitations

The study had some limitations. The first limitation was a small sample size. The second limitation is excluding from the study patients having GFR <60 mL/min/1.73 m². The third limitation is a result of an absence of the subpopulation of normotensive patients having T2DM. However, these limitations do not lead to a significant misunderstanding of the results received in the study.

Conclusion:

Cardiac remodeling in hypertensive patients having T2DM was significantly associated with higher levels of sST2 and cystatin C. sST2 had the best potency to predict LV remodeling when compared with NGAL and other biomarkers.

Authors' contribution

OMB designed the study protocol and proposed the idea of investigation. OMB, IMK, MVB, DAK, and OA obtained findings, analyzed the data, and reviewed the manuscript. AEB analyzed the data and wrote, edited, and reviewed the manuscript. MVB and LPA performed data collection, analyzed the statistics data, and checked the accuracy of the data analysis. IMK, LPA, and OMK performed ultrasound examinations and analyzed the results of it. OMK and LPA collected blood samples and performed the measurement of circulating biomarker concentrations. All authors read and approved the final version of the manuscript.

Acknowledgments

We thank staff person of Clinical Pharmacology and Internal Medicine department (Kharkiv National Medical University, Kharkiv, Ukraine) and Government Institution "L.T.Malaya Therapy National Institute NAMSU" (Kharkiv, Ukraine) as well as patients for participation in the study. We also thank a team of foreign languages department (Kharkiv National Medical University, Kharkiv, Ukraine) for assistance in manuscript preparation.

Competing interests

Authors have no competing interests

Ethics approval and consent to participate

The study was approved by the local ethics committee of Kharkiv National Medical University, Kharkiv, Ukraine. All patients gave their voluntary consent to participate in the study

Funding

This study was not supported by the non-commercial organizations and funds. Kharkiv National Medical University was the only source of financial supporting.

Abbreviations

ACE – angiotensin-converting enzyme
ADA – American Diabetic Association
ASE - American Society of Echocardiography
BMI - body mass index
CHF – chronic heart failure
CV – cardiovascular
DBP - diastolic blood pressure
ECS – European Cardiology Society
GFR - glomerular filtration rate
HbA_{1c} – glycosylated hemoglobin
HDL - high-density lipoprotein cholesterol
HOMA-IR – Homeostatic Assessment Model of Insulin Resistance
HR – heart rate

LDL - low-density lipoprotein cholesterol
 LV – left ventricular
 LVMM - LV myocardial mass
 LVMMI – LV myocardial mass index
 SBP – systolic blood pressure
 T2DM – type 2 diabetes mellitus.

References

1. Niazi M, Yari F, Shakarami A. A Review of Medicinal Herbs in the Lamiaceae Family Used to Treat Arterial Hypertension. *Entomol. Appl. Sci. Lett.* 2019 May 4;6(1):22-7.
2. Alshali KZ. Review of herb supplement use in type 2 diabetes. *Arch. Pharm. Pract.* 2020 Apr 1;11(2):42-9.
3. Krishnamoorthy Y, Sakthivel M, Sarveswaran G, Eliyas SK. Effectiveness of peer led intervention in improvement of clinical outcomes among diabetes mellitus and hypertension patients-A systematic review and meta-analysis. *Prim Care Diabetes.* 2019; 13(2):158-169. doi: 10.1016/j.pcd.2018.11.007.
4. Dunlay SM, Givertz MM, Aguilar D, Allen LA, Chan M, Desai AS, Deswal A, Dickson VV, Kosiborod MN, Lekavich CL, McCoy RG. American Heart Association Heart Failure and Transplantation Committee of the Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; and the Heart Failure Society of America. Type 2 Diabetes Mellitus and Heart Failure: A Scientific Statement from the American Heart Association and the Heart Failure Society of America: This statement does not represent an update of the 2017 ACC/AHA/HFSA heart failure guideline update. *Circulation.* 2019; 140(7):e294-e324. doi: 10.1161/CIR.0000000000000691.
5. Cryer MJ, Horani T, DiPette DJ. Diabetes and Hypertension: A Comparative Review of Current Guidelines. *J Clin Hypertens (Greenwich).* 2016;18(2):95-100. doi: 10.1111/jch.12638.
6. Petrie JR, Guzik TJ, Touyz RM. Diabetes, Hypertension, and Cardiovascular Disease: Clinical Insights and Vascular Mechanisms. *Can J Cardiol.* 2018; 34(5):575-584. doi: 10.1016/j.cjca.2017.12.005.
7. Berezin AE, Berezin AA, Lichtenauer M. Emerging Role of Adipocyte Dysfunction in Inducing Heart Failure Among Obese Patients With Prediabetes and Known Diabetes Mellitus. *Front. Cardiovasc. Med.* 2020; 7:583175. doi: 10.3389/fcvm.2020.583175.
8. Tan LM, Bakry MM, Redzuan A. Physiological Response to Dietary Salt: A Review on Salt Sensitivity Hypertension. *Int. J. Pharm. Phytopharm. Res.* 2018 Apr 1;8(2):8-20.
9. Alismaail AI, Alosaimi WA, Faqih AY, Al-Sahagi RJ, Alfraiji AF, Aloofy OA, Alhaidar SM, Alfaifi AA, Alrasheed MS, Alkhurayb NT. Hypertension Diagnosis and Management Approach. *Int. J. Pharm. Res. Allied Sci.* 2020 Jan 1;9(1):84-8.
10. Gan WZ, Ramachandran V, Lim CSY, Koh RY. Omics-based biomarkers in the diagnosis of diabetes. *J Basic Clin Physiol Pharmacol.* 2019; 31(2) doi: 10.1515/jbpcpp-2019-0120. PMID: 31730525.
11. Berezin A. Cardiac biomarkers in diabetes mellitus: New dawn for risk stratification? *Diabetes Metab Syndr.* 2017;11 Suppl 1:S201-S208. doi: 10.1016/j.dsx.2016.12.032
12. Gyöngyösi M, Winkler J, Ramos I, Do QT, Firat H, McDonald K, González A, Thum T, Díez J, Jaisser F, Pizard A. Myocardial fibrosis: biomedical research from bench to bedside. *Eur J Heart Fail.* 2017; 19(2): 177–191. doi: 10.1002/ejhf.696
13. Colhoun HM, Marcovecchio ML. Biomarkers of diabetic kidney disease. *Diabetologia.* 2018;61(5):996-1011. doi: 10.1007/s00125-018-4567-5.
14. Bozkurt B, Aguilar D, Deswal A, Dunbar SB, Francis GS, Horwich T, Jessup M, Kosiborod M, Pritchett AM, Ramasubbu K, Rosendorff C, Yancy C; American Heart Association Heart Failure and Transplantation Committee of the Council on Clinical Cardiology; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular and Stroke Nursing; Council on Hypertension; and Council on Quality and Outcomes Research. Contributory Risk and Management of Comorbidities of Hypertension, Obesity, Diabetes Mellitus, Hyperlipidemia, and Metabolic Syndrome in Chronic Heart Failure: A Scientific Statement From the American Heart Association. *Circulation.* 2016; 134(23):e535-e578. doi: 10.1161/CIR.0000000000000450.
15. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, De Simone G, Dominiczak A, Kahan T. 2018 ESC/ESH Guidelines for the management of arterial hypertension. *Eur Heart J.* 2018; 39(33): 3021–3104
16. De Boer IH, Bangalore S, Benetos A, Davis AM, Michos ED, Muntner P, Rossing P, Zoungas S, Bakris G. Diabetes and hypertension: a position statement by the American Diabetes Association. *Diabetes Care.* 2017 Sep 1;40(9):1273-84.
17. Marwick TH, Gillebert TC, Aurigemma G, Chirinos J, Derumeaux G, Galderisi M, Gottdiener J, Haluska B, Ofili E, Segers P, Senior R. Recommendations on the Use of Echocardiography in Adult Hypertension: A Report from the European Association of Cardiovascular Imaging (EACVI) and the American Society of Echocardiography (ASE). *J Am Soc Echocardiogr.* 2015; 28(7):727-54. doi: 10.1016/j.echo.2015.05.002
18. Nagueh SF, Smiseth OA, Appleton CP, Byrd BF 3rd, Dokainish H, Edvardsen T, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of

- Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr.* 2016; 29(4):277-314. doi: 10.1016/j.echo.2016.01.011.
19. Levey AS, Stevens LA, Schmid CH, Zhang Y, Castro III AF, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J. for the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A New Equation to Estimate Glomerular Filtration Rate. *Ann Intern Med.* 2009; 150(9): 604-12
 20. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985; 28(7):412-9
 21. Gilca GE, Stefanescu G, Badulescu O, Tanase DM, Bararu I, Ciocoiu M.. Potential Diagnostic and Therapeutic Targets. *J Diabetes Res.* 2017; 2017:1310265. doi: 10.1155/2017/1310265
 22. Lee WS., Kim J. Diabetic cardiomyopathy: where we are and where we are going. *Korean J Intern Med.* 2017;32(3):404-421. doi: 10.3904/kjim.2016.208
 23. Anders HJ, Huber TB, Isermann B, Schiffer M. CKD in diabetes: diabetic kidney disease versus nondiabetic kidney disease. *Nat Rev Nephrol.* 2018;14(6):361-377. doi: 10.1038/s41581-018-0001-y.
 24. Koye DN, Magliano DJ, Reid CM, Jepson C, Feldman HI, Herman WH, Shaw JE. Risk of Progression of Nonalbuminuric CKD to End-Stage Kidney Disease in People With Diabetes: The CRIC (Chronic Renal Insufficiency Cohort) Study. *Am J Kidney Dis.* 2018; 72(5):653-661. doi: 10.1053/j.ajkd.2018.02.364. Epub 2018 May 18. PMID: 29784612.
 25. Khangura KK, Eirin A., Kane GC., Misra S., Textor S., Lerman A., Lerman LO. Cardiac function in renovascular hypertensive patients with and without renal dysfunction. *Am J Hypertens.* 2014; 27(3):445-53. doi: 10.1093/ajh/hpt203
 26. van Veldhuisen DJ., Ruilope LM., Maisel AS., Damman K Biomarkers of renal injury and function: diagnostic, prognostic and therapeutic implications in heart failure. *Eur. Heart J.* 2016; 37(33):2577-85. doi: 10.1093/eurheartj/ehv588
 27. Matys U, Bachorzewska-Gajewska H, Malyszko J, Dobrzycki S. Assessment of kidney function in diabetic patients. Is there a role for new biomarkers NGAL, cystatin C and KIM-1? *Adv Med Sci.* 2013; 58(2):353-61. doi: 10.2478/v10039-012-0077-8.
 28. Bansal N, Zelnick L, Shlipak MG, Anderson A, Christenson R, Deo R, Defilippi C, Feldman H, Lash J, He J, Kusek J. CRIC Study Investigators. Cardiac and Stress Biomarkers and Chronic Kidney Disease Progression: The CRIC Study. *Clin Chem.* 2019; 65(11):1448-1457. doi: 10.1373/clinchem.2019.305797.
 29. Alicic RZ, Rooney MT, Tuttle KR. Diabetic Kidney Disease: Challenges, Progress, and Possibilities. *Clin J Am Soc Nephrol.* 2017; 12(12):2032-2045. doi: 10.2215/CJN.11491116
 30. Piéroni L, Cristol JP. Biomarqueurs urinaires de dysfonction rénale [Urinary biomarkers of kidney dysfunction]. *Ann Biol Clin (Paris).* 2015; 73(2):151-7. doi: 10.1684/abc.2015.1029.
 31. Ofstad AP, Atar D, Gullestad L, Langslet G, Johansen OE. The heart failure burden of type 2 diabetes mellitus-a review of pathophysiology and interventions. *Heart Fail Rev.* 2018; 23(3): 303-323. doi: 10.1007/s10741-018-9685-0..
 32. Gibb AA, Hill BG. Metabolic Coordination of Physiological and Pathological Cardiac Remodeling. *Circ Res.* 2018; 123(1):107-128. doi: 10.1161/CIRCRESAHA.118.312017. PMID: 29929976; PMCID: PMC6023588.
 33. Jia G, Hill MA, Sowers JR. Diabetic Cardiomyopathy: An Update of Mechanisms Contributing to This Clinical Entity. *Circ Res.* 2018; 122(4): 624-638. doi: 10.1161/CIRCRESAHA.117.311586.
 34. Bertero E, Maack C. Metabolic remodelling in heart failure. *Nat Rev Cardiol.* 2018; 15(8):457-470. doi: 10.1038/s41569-018-0044-6.
 35. Yap J, Tay WT, Teng TK, Anand I, Richards AM, Ling LH, et al; ASIAN-HF (Asian Sudden Cardiac Death in Heart Failure) Registry Investigators, Zile M, McMurray J, Lam CSP. Association of Diabetes Mellitus on Cardiac Remodeling, Quality of Life, and Clinical Outcomes in Heart Failure With Reduced and Preserved Ejection Fraction. *J Am Heart Assoc.* 2019; 8(17):e013114. doi: 10.1161/JAHA.119.013114.
 36. Gonçalves N, Gomes-Ferreira C, Moura C, Roncon-Albuquerque R Jr, Leite-Moreira AF, Falcão-Pires I. Worse cardiac remodeling in response to pressure overload in type 2 diabetes mellitus. *Int J Cardiol.* 2016; 217:195-204. doi: 10.1016/j.ijcard.2016.04.178.