

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/373103001>

The Theoretical Description for Ibotenic Acid and Muscimol Electrochemical Determination in Mushroom Pulp and Mushroom-based Alcoholic Beverages on Nano- CuS Composite with Conduct...

Article in *Letters in Applied NanoBioScience* - March 2024

DOI: 10.33263/LIANBS131.037

CITATIONS

0

READS

31

27 authors, including:



Volodymyr Tkach

Chernivtsi National University

714 PUBLICATIONS 473 CITATIONS

[SEE PROFILE](#)



Marta Kushnir

Chernivtsi National University

426 PUBLICATIONS 125 CITATIONS

[SEE PROFILE](#)



Lyudmyla Romaniv

Chernivtsi National University

67 PUBLICATIONS 34 CITATIONS

[SEE PROFILE](#)



Silvio de Oliveira

Universidade Federal de Mato Grosso do Sul

488 PUBLICATIONS 1,011 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



potentiometric sensors based on ion associate with base dye [View project](#)



wildlife corridor, habitat corridor, green corridor [View project](#)

The Theoretical Description for Ibotenic Acid and Muscimol Electrochemical Determination in Mushroom Pulp and Mushroom-based Alcoholic Beverages on Nano-CuS Composite with Conducting Polymer

Volodymyr V. Tkach ^{1,*}, Marta V. Kushnir ¹, Lyudmyla V. Romaniv ¹, Sílvio C. de Oliveira ², Yana G. Ivanushko ³, Yevgeniya V. Nazymok ³, Oleksandra V. Ahafonova ³, Petro I. Yagodynets ^{1,*}, Adriano O. da Silva ⁴, Nataliia P. Derevianko ⁵, Mykhailo P. Zavorodnii ⁵, Vira M. Odyntsova ⁶, Mykola P. Krasko ⁶, Olena O. Vasylieva ⁷, Olena V. Hrabovska ⁷, Dilfuza M. Musayeva ⁸, Riboba Sharipova ⁸, Bakhodirjon Samadov ⁸, Laziz N. Niyazov ⁸, Tetiana V. Morozova ⁹, Jarem R. Garcia ¹⁰, José Inácio Ferrão da Paiva Martins ¹¹, Zholt O. Kormosh ¹², Alla V. Grekova ¹³, Ianina F. Burdina ¹³, Inesa M. Khmel'iar ¹⁴, Lesya O. Kushnir ¹⁴

¹ Chernivtsi National University, 58001, Kotsyubynsky Str. 2, Chernivtsi, Ukraine

² Institute of Chemistry. Federal University of Mato Grosso do Sul, 79074 – 460, Av. Sen. Felinto Müller, 1555, Vila Ipiranga, Campo Grande, MS, Brazil

³ Bukovinian State Medical University, 58001, TeatralnaSq, 9, Chernivtsi, Ukraine

⁴ Federal University of the West of Pará, Juruti Campus, 68170 – 000, Rua Veríssimo de Souza Andrade, s/n, Juruti, PA, Brazil

⁵ Khortytsia National Rehabilitation Academy, 69000, Naukove Mistechko, 59, Khortytsia Island, Zaporizhzhia, Ukraine

⁶ Zaporizhzhia State Medical University, 69600, Mayakovsky Ave. 24, Zaporizhzhia, Ukraine

⁷ Kyiv National University of Trade and Economics, 02156, Kyoto Str. 21, Kyiv, Ukraine~

⁸ Abu Ali Ibn Sino Bukhara State Medical Institute, 705018, Navoi Str., 1, Bukhara, Uzbekistan

⁹ National University of Food Technologies, Volodymyrska Str., 68, 01001, Kyiv, Ukraine

¹⁰ State University of Ponta Grossa, Uvaranas Campus, Av. Gal. Carlos Cavalcanti, 4748, 84030-900, Ponta Grossa, PR, Brazil

¹¹ Engineering Faculty of the University of Porto, 4200-465, Rua Dr. Roberto Frias, s/n, Porto, Portugal

¹² Volyn National University, 43000, Voli Ave., 13, Lutsk, Ukraine

¹³ Odesa National Medical University, 65000, Valikhovsky Ln. 2, Odesa, Ukraine

¹⁴ Rivne State Basic Medical Academy, 33000, Mykola Karnaukhov Str., 53, Rivne, Ukraine

* Correspondence: nightwatcher2401@gmail.com (V.V.T.), ved1988mid@rambler.ru (P. I.Y.);

Scopus Author ID 55758299100

Received: 27.02.2023; Accepted: 31.03.2023; Published: 13.08.2023

Abstract: In this article, the possibility for ibotenic acid and muscimol electrochemical determination on the electrode, modified by nano-CuS/conducting polymer composite, is theoretically described. The analytical signal is based on the reaction of the copper (III) sulfohydroxide with both of the analytes, and the reaction is realized by two parallel scenarios. The analysis of the correspondent model confirms the possibility of the oscillatory behavior caused by DEL influence of the electrochemical stage and chemical stage involving ibotenic acid. The chemical stages involving muscimol do not participate in the oscillatory behavior. The stability analysis confirms the easy realization of the linear dependence between the electrochemical parameter and concentration and, therefore, the facility in analytical signal interpretation.

Keywords: *Amanita muscaria*; ibotenic acid; muscimol; electrochemical sensor; conducting polymer; copper (II) sulfide; stable steady-state.

© 2023 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Fly agaric, also known as *Amanita muscaria* [1–4], is one of the most widespread poisonous mushrooms in subtropical and moderate countries. It also becomes widespread in subequatorial and equatorial countries such as Panama, Colombia, and Venezuela [4]. Its poison is highly toxic, causing grave central nervous system disorders [5]. Nevertheless, it is still used in some alcoholic drinks in Russia, Finland, and Norway [6,7]. In Middle Ages, the vikings ate fly agaric into mushroom broth in order to enhance the spirit.

The fly agaric toxicity is generally caused by the presence of ibotenic acid [8–11] and their derivatives – muscimol and muscazone in the mushroom pulp (Figure 1).




Figure 1. Ibotenic acid, muscimol and muscazone.

Being muscimol is the most dangerous among the three [12–14]. Moreover, being a biogenic amine, it stimulates the nervous system, causing the false happiness feeling and giving the reforce to the action of two other toxins. As for muscazone, a product of ibotenic acid photoisomerization, its action is less intensive but longer and somehow more intense (including partial blindness due to the methanol formation in its metabolism).

Considering the structure of all three compounds, we may conclude that cathodic reduction is the best strategy to detect all of the compounds electrochemically. Nevertheless, anodic oxidation may also be used. Muscimol will be oxidized at lower potentials than ibotenic acid and muscazone. For this purpose, chemically modified electrodes containing strong oxidants may be used [15–18].

Therefore, the composite material of CuS nanoparticles with a conducting polymer, in which the polymer is a mediator and stabilizer, and the CuS is the starting material for *in situ* anodic synthesis of trivalent copper as copper (III) sulfohydroxide by reaction (1):



would be an interesting option. Such hybrid materials have already been used in electroanalytic[19–22] in either cathodic or anodic processes. Nevertheless, the electrochemical instabilities, typical for similar electroanalytical and electro synthetical processes [22–26], may impede the easy interpretation of the analytical signal or even lead to the electrochemical equipment breaking down. On the other hand, the electrochemical instabilities may reveal the presence of a substance or a specific reaction involving the electroactive compound.

The analysis of the theoretical and experimental data shows that the electrochemical instabilities in non-autocatalytic electrochemical systems are caused by the influences of

chemical and electrochemical processes on the double electric layer (DEL) ionic force, capacity, and conductivity, like also the electrode surface material impedance.

Therefore, the theoretical *a priori* investigation of the system's behavior, capable of detecting the conditions of the most effective sensing interpretation, like also the conditions of the electrochemical instabilities, is an important stage for the development of new electrochemical sensors, especially for new compounds and involving new electrode modifiers. Also, this investigation, involving the development and analysis of a mathematical model, permits to compare the similar electroanalytical systems. Therefore, the aim of this work is to investigate, from the mechanistic theoretical point of view, the possibility of ibotenic acid and muscimol electrochemical determination on nano-CuS/conducting polymer composite. It involves the mechanism suggestion and mathematical model analysis and interpretation, including the comparison of this system's behavior with that of similar ones [27,28].

2. Materials and Methods

Copper (III) sulfohydroxide, a product of copper (II) sulfide *in situ* electrooxidation by reaction (1), is an aggressive oxidant, oxidizing both ibotenic acid and muscimol by two parallel scenarios:

- yielding an N-oxide;
- oxidizing amino group to imine.

Considering the absence of highly accepting carboxyl in muscimol, its oxidation will be given more easily and on lower anodic potential values (Figure 2).




Figure 2. The scheme for the electroanalytical process.

Taking this into account and taking some assumptions [27,28], we describe the behavior of this system by a trivariant balance differential equation-set (2):

$$\begin{cases} \frac{dm_1}{dt} = \frac{2}{\delta} \left(\frac{M_1}{\delta} (m_{10} - m_1) - r_{11} - r_{12} \right) \\ \frac{dm_2}{dt} = \frac{2}{\delta} \left(\frac{M_2}{\delta} (m_{20} - m_2) - r_{21} - r_{22} \right) \\ \frac{dc}{dt} = \frac{1}{c} (r_{11} + r_{12} + r_{21} + r_{22} - r_1) \end{cases} \quad (2)$$

Herein, m_1 and m_2 are correspondently ibotenic acid and muscimol concentrations in the pre-surface layer, δ is the diffusion thickness, c is the copper sulfide matrix coverage degree, M_1 and M_2 are ibotenic acids and muscimol diffusion coefficients, m_{10} and m_{20} stand for the bulk concentration of each of the analytes, C is the copper sulfide maximal matrix concentration, and the parameters r stand for the correspondent reaction rates, calculated as:

$$r_{11} = k_{11}m_1(1 - c)^2 \exp(-ac) \quad (3)$$

$$r_{12} = k_{12}m_1(1 - c)^2 \exp(-ac) \quad (4)$$

$$r_{21} = k_{21}m_2(1 - c)^2 \quad (5)$$

$$r_{22} = k_{22}m_2(1 - c)^2 \quad (6)$$

$$r_1 = k_1c \exp\left(\frac{F\varphi_0}{RT}\right) \quad (7)$$

Herein the parameters k correspond to the reaction rate constants, a is the parameter describing the influence of ionic forms transformations on DEL electrophysical properties, $F = N_Ae$ is the Faraday constant, φ_0 stands for zero-charge-related potential slope in DEL, R is the universal gas constant, and T is the absolute temperature.

In this system, the oscillatory behavior on the chemical stage may be caused by the DEL impact of only one of two analytes analyte. Therefore, the electroanalytical process will be more stable and efficient than for ibotenic acid and muscazone, as shown below.

3. Results and Discussion

We investigate the behavior of the system with ibotenic acid and muscimol determination over CuS/conducting polymer composite by means of the linear stability theory, describing the steady-state Jacobian members as (8):

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad (8)$$

In which:

$$a_{11} = \frac{2}{\delta} \left(-\frac{M_1}{\delta} - k_{11}(1 - c)^2 \exp(-ac) + ak_{11}m_1(1 - c)^2 \exp(-ac) - k_{12}(1 - c)^2 \exp(-ac) + ak_{12}m_1(1 - c)^2 \exp(-ac) \right) \quad (9)$$

$$a_{12} = 0 \quad (10)$$

$$a_{13} = \frac{2}{\delta} (2k_{12}m_1(1 - c) \exp(-ac) + 2k_{12}m_1(1 - c) \exp(-ac)) \quad (11)$$

$$a_{21} = 0 \quad (12)$$

$$a_{22} = \frac{2}{\delta} \left(-\frac{M_2}{\delta} - k_{21}(1 - c)^2 - k_{22}(1 - c)^2 \right) \quad (13)$$

$$a_{23} = \frac{2}{\delta} (2k_{21}m_2(1 - c) + 2k_{22}m_2(1 - c)) \quad (14)$$

$$a_{31} = \frac{1}{c} (k_{11}(1 - c)^2 \exp(-ac) - ak_{11}m_1(1 - c)^2 \exp(-ac) + k_{12}(1 - c)^2 \exp(-ac) - k_{12}m_1(1 - c)^2 \exp(-ac)) \quad (15)$$

$$a_{32} = \frac{1}{c} (k_{21}(1 - c)^2 + k_{22}(1 - c)^2) \quad (16)$$

$$a_{33} = \frac{1}{c} \left(-2k_{12}m_1(1 - c) \exp(-ac) - 2k_{12}m_1(1 - c) \exp(-ac) - 2k_{21}m_2(1 - c) - 2k_{22}m_2(1 - c) - k_1 \exp\left(\frac{nF\phi_0}{RT}\right) + jk_1c \exp\left(\frac{nF\phi_0}{RT}\right) \right) \quad (17)$$

Taking into account the Jacobian main diagonal elements (6), (10), and (14), it is possible to conclude that, as this diagonal contains the elements capable of being positive, the positive callback may be characteristic of this electrochemical process. Therefore, the Hopf bifurcation, correspondent to the oscillatory behavior, becomes possible.

In this system, the oscillatory behavior is caused by the (co)action of two factors, such as in [27 - 28]. Besides the factor of the influence of the electrochemical stage on the surface and pre-surface layer electrophysical properties (ionic force, conductivity, and capacitance), described by the positivity of the element $jk_1c \exp\left(\frac{nF\phi_0}{RT}\right) > 0$, if $j > 0$, the oscillatory behavior will also be caused by ionic form transformation on the chemical stage, described by the positivity of the elements $ak_{11}m_1(1 - c)^2 \exp(-ac)$ and $ak_{12}m_1(1 - c)^2 \exp(-ac)$, involving the reactions of the ibotenate-ion.

It's important to mention that in neutral and mildly alkaline mediums, the second analyte, muscimol, isn't ionized. Therefore, in this system, the oscillatory behavior will be defined by the chemical transformation of only one analyte – ibotenic acid in the form of ibotenate-ion.

As for the steady-state stability, its topological region is more narrow than in the simplest case but wider than in the case of two ionizing analytes. Applying the Routh-Hurwitz criterion, we rewrite the Jacobian matrix as (18):

$$\frac{4}{\delta^2 C} \begin{vmatrix} -\kappa - E & 0 & \Sigma \\ 0 & -\lambda - \Lambda & T \\ E & \Lambda & -\Sigma - T - \Omega \end{vmatrix} \quad (18)$$

Introducing new variables in order to avoid cumbersome expressions.

Opening the straight brackets and resolving the inequation $\text{Det } J < 0$, salient from the criterion, and changing the signs to the opposite, we obtain the necessary requisite for steady-state stability (19):

$$\kappa(\lambda\Sigma + \lambda T + \lambda\Omega + \Lambda\Sigma + \Lambda\Omega) + E(\lambda T + \lambda\Omega + \Lambda\Omega) > 0 \quad (19)$$

This inequity describes an efficient electroanalytical process, which is both diffusion and kinetically controlled. Moreover, taking into account the absence of side reactions, compromising the analyte and modifier stability, it will depend on linear dependence between the analytes concentration and electrochemical parameter (current and/or) potential).

As for the detection limit, it will correspond to the monotonic instability, depicting the margin between the stable steady-state and unstable state. Its condition may be exposed as $\text{Det } J = 0$ or (19):

$$\kappa(\lambda\Sigma + \lambda T + \lambda\Omega + \Lambda\Sigma + \Lambda\Omega) + E(\lambda T + \lambda\Omega + \Lambda\Omega) = 0 \quad (20)$$

A similar scenario will be given in the case of muscimol and muscazone electrochemical determination. Nevertheless, it will be simplified for muscazone due to only one oxidation possibility against two for muscimol and ibotenic acid. This behavior is

characteristic till a certain pH value, at which the amido ester group in muscazone is hydrolyzed. The behavior of this system will be described in one of our next works.

4. Conclusions

From the analysis of the system with ibotenic acid and muscimol determination on CuS/Conducting Polymer composite, it is possible to conclude that the mentioned composite is an efficient electrode modifier for ibotenic acid and muscimol determination. The electroanalytical process is both diffusion and kinetically controlled, being less stable than for the simplest case but more stable than for two ionizing analytes. As for the oscillatory behavior, its probability is high but lower than for the case of ibotenic acid and muscazone.

Funding

This research received no external funding.

Acknowledgments

Volodymyr V. Tkach acknowledges the Engineering Faculty of the University of Porto and the University of Trás-os-Montes and Alto Douro for their support during these difficult times for Ukraine and its research.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Rampoll, F.I.; Kamler, P.; Carlino, C.C.; Bedussi, F. The Deceptive Mushroom: Accidental *Amanita muscaria* Poisoning. *Eur. J. Case Rep. Int. Med.* **2021**, *8*, https://doi.org/10.12890/2021_002212.
2. Carboue, Q.; López, M. *Amanita muscaria*: Ecology, Chemistry, Myths. *Encyclopedia* **2021**, *1*, 905–914, <https://doi.org/10.3390/encyclopedia1030069>.
3. Ittiyavirah, S.P.; Ramalingam, K.; Sathyan, A.; Rajasree, R.S.; Kuruniyan, M.S.; Quadri, S.A.; Elayadeth-Meethal, M.; Naseef, P.P. Thymoquinone-Rich Black Cumin Oilk Attenuates Ibotenic Acid Induced Excitotoxicity Through Glutamate Receptors in Wistar Rats. *Saudi Pharm J.* **2022**, *30*, 1781–1790, <https://doi.org/10.1016/j.jsps.2022.10.007>.
4. Vargas, N.; Gonçalves, S.C.; Franco-Molano, A.E.; Restrepo, S.; Pringle, A. In Colombia the Eurasian fungus *Amanita muscaria* is expanding its range into native, tropical *Quercus humboldtii* forests. *Mycologia* **2019**, *111*, 758–771, <https://doi.org/10.1080/00275514.2019.1636608>.
5. Gao, J.; Xie, R.; Wang, N.; Zhang, J.; Sun, X.; Wang, H.; Tan, J.; Chen, A. Rapid identification of *Amanita citrinoannulata* poisoning using colorimetric and real-time fluorescence and loop-mediated isothermal amplification (LAMP) based on the nuclear ITS region. *Food Chemistry: Molecular Sciences* **2022**, *4*, <https://doi.org/10.1016/j.fochms.2022.100082>.
6. Winkelman, M.J. *Amanita muscaria*: Fly Agaric History, Mythology and Pharmacology. *J. Psychedel. Stud.* **2022**, *6*, 1–4, <https://doi.org/10.1556/2054.2022.00216>.
7. Housecroft, C.E. The fungus *Amanita Muscaria*. From Neutoroxin to Vanadium Accumulation. *Chimia* **2019**, *73*, 96–97, <https://doi.org/10.2533/chimia.2019.96>.
8. Modaberi, S.; Shahbazi, M.; Naghdi, N.; Bagherzadeh, F. Effects of bilateral microinjections of ibotenic acid into neostriatal region and forced exercise on spatial learning and memory and anthropometric characteristics of male rats. *Koomesh* **2021**, *23*, 654–664, <https://doi.org/10.52547/koomesh.23.5.654>.
9. Rostampour, M.; Tehrani, B.S.; Babaei, P. Effect of ibotenic acid and colchicine infusion into the rat hippocampus in passive avoidance learning. *Pathobiol. Res.* **2021**, *24*, 37–43.
10. Li, H.; Zhang, H.; Zhang, Y.; Zhang, K.; Zhou, J.; Yin, Y.; Jiang, S.; Ma, P.; He, Q.; Zhang, Y. Mushroom poisoning outbreaks—China, 2019. *China CDC Weekly* **2020**, *2*, <https://doi.org/10.46234/ccdcw2020.005>.
11. Sasaki, T.; Saito, H.; Hiradate, Y.; Hara, K.; Tanemura, K. Behavioural effects in mice orally exposed to domoic acid or ibotenic acid are influenced by developmental stages and sex differences. *Biochemical and Biophysical Research Communications* **2021**, *558*, 175–182, <https://doi.org/10.1016/j.bbrc.2021.04.080>.

12. Lemerrier, C.E.; Garenne, A.; Poullietier de Gannes, F.; El Khoueiry, C.; Arnaud-Cormos, D.; Levêque, P.; Lagroye, I.; Percherancier, Y.; Lewis, N. Comparative Study Between Radiofrequency-Induced and Muscimol-Induced Inhibition of Cultured Networks of Cortical Neuron. *PLoS One* **2022**, *17* <https://doi.org/10.1371/journal.pone.0268605>.
13. Kim, E.-J.; Kwon, O.-S.; Hur, C.-G.; Nyiramana, M.M.; Lee, D.-K.; Hong, S.-G.; Han, J.; Kang, D. Muscimol Directly Activates the TREK-2 Channel Expressed in GABAergic Neurons through Its N-Terminus. *International Journal of Molecular Sciences* **2021**, *22*, <https://doi.org/10.3390/ijms22179320>.
14. Sawangjit, A.; Harkotte, M.; Oyanedel, C.N.; Niethard, N.; Born, J.; Inostroza, M. Two Distinct Ways to Form Long-Term Object Recognition Memory During Sleep and Wakefulness. *Psych. Cogn. Sci. Neuroscience* **2022**, *119*, e2203165119, <https://doi.org/10.1073.pnas.2203165119>.
15. Obermaier, S.; Müller, M. Ibotenic Acid Biosynthesis in the Fly Agaric is Initiated by Glutamate Hydroxylation. *Angew. Chem. Int. Ed.* **2020**, *59*, 12432–12435, <https://doi.org/10.1002/anie.202001870>.
16. Lee, S.; Yu, Y.S.; Lee, S.R.; Kim, K.H. Non-peptide secondary metabolites from poisonous mushrooms: overview of chemistry, bioactivity, and biosynthesis. *Nat. Prod. Rep.* **2022**, *39*, 512–559, <https://doi.org/10.1039/D1NP00049G>.
17. Poliwoda, A.; Zielinska, K.; Halama, M.; Wieczorek, P. Determination of muscimol and ibotenic acid in mushrooms of Amanitaceae by capillary electrophoresis. *Electrophoresis*, **2014**, *35*, <http://dx.doi.org/10.1002/elps.201400104>.
18. Joshi, N.C.; Malik, S.; Gururani, P. Utilization of Polypyrrole/ZnO Nanocomposite in the Adsorptive Removal of Cu²⁺, Pb²⁺ and Cd²⁺ Ions from Wastewater. *Letters in Applied NanoBioScience* **2021**, *10*, 2339–2351, <https://doi.org/10.33263/LIANBS103.23392351>.
19. Santos de Oliveira, R.W.; Barros Gomes, P.R.; Ribeiro, J.D.; Batista Reis, J.; Alves Fontenele, M.; Oliveira Everton, G.; da Silva Lyra, W.; Costa Louzeiro, H.; Livramento de Paula, M.d.; Mouchrek Filho, V.E. Determinação Espectrofotométrica de Cobre (II) em Aguardente de Mandioca (Tiquira) (in Portuguese). *Rev. Colomb. Cien. Quím. Farm.* **2020**, *49*, 353–371, <https://doi.org/10.15446/rcciquifa.v49n2.89507>.
20. Debbarma, J.; Debnath, R.; Saha, M. Graphene from Sugarcane Bagasse for Nonenzymatic Electrochemical Determination of Glucose. *Lett. Appl. NanoBioSci.* **2023**, *10*, <https://doi.org/10.33263/LIANBS124.115>.
21. Zou, X.; Deng, Z.; Chen, H.; Zheng, Z.; Ji, L.; Chen, Y.; Sun, M.; Ouyang, S.; Yuan, Z.; Zhao, P.; Tao, J. Dual-Signal Colorimetric and Electrochemical Sensor of Dopamine Based on Nanocomposite of Cobalt Oxyhydroxide/Carbon Black. *J. Electrochem. Soc.* **2023**, *170*, <https://doi.org/10.1149/1945-7111/acb237>.
22. Bialas, K.; Moschou, D.; Marken F.; Estrela, P. Electrochemical Sensors Based on Metal Nanoparticles with Biocatalytic Activity. *Microchim. Acta.* **2022**, *189*, <https://doi.org/10.1007/s00604-022-05252-2>.
23. Holze, R. Overoxidation of Intrinsically Conducting Polymers. *Polymers* **2022**, *14*, <https://doi.org/10.3390/polym14081584>.
24. Thadathil, A.; Pradeep, H.; Joshy, D.; Ismail, Y.A.; Periyat, P. Polyindole and Polypyrrole as a Sustainable Platform for Environmental Remediation and Sensor Applications. *Mater Adv.* **2022**, *3*, 2990–3022, <https://doi.org/10.1039/D2MA00022A>.
25. Das, I.; Goel, N.; Agrawal, N.R.; Gupta, S.K. Growth Patterns of Dendrimers and Electric Potential Oscillations during Electropolymerization of Pyrrole using Mono- and Mixed Surfactants. *The Journal of Physical Chemistry B* **2010**, *114*, 12888–12896, <https://doi.org/10.1021/jp105183q>.
26. Das, I.; Goel, N.; Gupta, S.K.; Agrawal, N.R. Electropolymerization of pyrrole: Dendrimers, nano-sized patterns and oscillations in potential in presence of aromatic and aliphatic surfactants. *Journal of Electroanalytical Chemistry* **2012**, *670*, 1–10, <https://doi.org/10.1016/j.jelechem.2012.01.023>.
27. Tkach, V.; Kushnir, M.; de Oliveira, S.; Salomova, H.; Jalilov, F.; Jalilova, F.; Musayeva, D.; Niyazov, L.; Ivanushko, Y.; Ahafonova, O.; Mytchenok, M.; Yagodynets, P.; Kormosh, Z.; Reis, L.; Palytsia, Y. The Theoretical Description for Fluoxetine Electrochemical Determination, Assisted by CoO(OH)-Nanoparticles, Deposited Over the Squaraine Dye. *Orbital - The Electronic Journal of Chemistry* **2021**, *13*, 53–57.
28. Tkach, V.V.; Kushnir, M.V.; Koptiika, V.V.; Luganska, O.V.; Omelyanchik, L.O.; Kormosh, Z.O.; Mytchenok, M.P.; Garcia, J.R.; Palamarek, K.V.; Bagrii, K.L. Theoretical Description for Ibotenic Acid and Muscazone Determination in Mushroom Pulp and Biological Liquids over Conducting Polymer-Modified Electrode. *Biointerface Res. Appl. Chem.* **2023**, *13*, <https://doi.org/10.33263/BRIAC133.275>.